

CASE REPORT

Open Access

Coexistence of HBsAg and HBsAb in a difficult-to-treat chronic hepatitis B: loss of HBsAg with entecavir plus tenofovir combination

Giovanni Galati¹, Antonio De Vincentis^{1*}, Umberto Vespasiani-Gentilucci¹, Paolo Gallo¹, Donatella Vincenti², Maria Carmela Solmone², Chiara Dell'Unto¹ and Antonio Picardi¹

Abstract

Background: Some reports have documented the coexistence of Hepatitis B surface Antigen (HBsAg) and anti-HBsAg antibodies (HBsAb) in patients with chronic hepatitis B (CHB), often in the absence of amino acid substitutions in the HBsAg sequences of the Hepatitis B Virus (HBV) genome able to explain an immunological escape variant. HBV genome has a very compact coding organization, with four partially overlapping open reading frames (ORFs). Because the reverse transcriptase region (rt) of HBV polymerase overlaps the HBsAg ORF, it is possible that some mutations in the HBsAg region correspond to mutations in the rt ORF, conferring resistance to current antiviral therapies.

This unique case explores the response to antiviral therapies of a CHB with concurrent HBsAg and HBsAb positivity, and analyse the clinical implications of possible mutations in rt and HBsAg ORFs.

Case presentation: Here we describe the case of a 59 year-old Italian man suffering from Hepatitis B envelope Antigen (HBeAg) positive CHB with concurrent HBsAb positivity. By ultra-deep pyro-sequencing (UDPS) technique, mutations conferring immunological escape or resistance to antiviral therapies were found neither in HBsAg nor in HBV rt ORFs, respectively.

The patient was unsuccessfully treated with interferon, adefovir monotherapy and adefovir plus entecavir combination. Surprisingly, during entecavir plus tenofovir combination, anti-HBe seroconversion and HBsAg loss were observed, while the titer of HBsAb persisted.

Conclusions: Concurrent HBsAg/HBsAb positivity in active CHB is a clinical and virological dilemma. In this setting, there are not consistent data about the response to conventional therapies and the immunological balance between host and virus remains so far unexplained. This is, to our knowledge, the first case described of a CHB with HBsAg/HBsAb positivity, wild type for clinically relevant mutations in HBsAg and rt ORFs, successfully treated with a combination of nucleos(t)ide analogues (NAs).

Keywords: HBeAg positive chronic hepatitis B, HBsAg, Anti-HBs, Coexistence, Ultra-deep pyro-sequencing, Immunological escape, Nucleos(t)ide analogues, Combination, Entecavir, Tenofovir

* Correspondence: a.devincentis@unicampus.it

¹Clinical Medicine and Hepatology Unit, Campus Bio Medico University of Rome, 200, Alvaro Del Portillo Street, Rome 00128, Italy
Full list of author information is available at the end of the article

Background

Hepatitis B Virus (HBV) can cause a self-limiting acute infection or a chronic hepatitis, depending on the interaction between the host's immune system and the virus.

Typically, the sign of HBV infection is the presence of Hepatitis B Surface Antigen (HBsAg) in the blood. On the other hand, the appearance of the neutralizing antibodies against HBsAg (HBsAb) usually indicates resolution of infection, both spontaneously and after therapy [1].

In this simple virological scenario, some reports have documented the coexistence of HBsAg and HBsAb in some patients with chronic hepatitis B (CHB), often in the absence of amino acid substitutions in the HBsAg sequence able to explain the escape of HBV from the HBsAb immune control [2,3]. HBV genome has a very compact coding organization, with four partially overlapping open reading frames (ORFs). Because the reverse transcriptase (rt) region of HBV polymerase overlaps the HBsAg ORF, it is possible that mutations in the HBsAg region correspond to mutations in the rt ORF, conferring resistance to nucleos(t)ide analogues (NAs) [4,5]. In addition, due to the quasispecies nature of the HBV genome in each infected individual, some mutations may be present in minor variants of viral population, being not detected by classical population sequencing. The powerful ultra-deep pyro-sequencing (UDPS) approach, based on next generation sequencing (NGS), has been recently used to obtain a complete description of HBV quasispecies, highlighting possible minor populations carrying mutations in the two overlapping ORFs [6].

This case is relevant for clinical virology because explores the response to antiviral therapies of a CHB with concurrent HBsAg and HBsAb positivity, in the absence of clinically relevant mutations in rt and HBsAg ORFs.

Case presentation


A 59-year-old Italian man was admitted on July 2006 to the Hepatology Unit of the University Hospital "Campus Bio-Medico" of Rome, for investigations concerning CHB. He had not been vaccinated against HBV, he had no known risk factors for contracting viral hepatitis, and all his households were negative for HBsAg. At the time of admission, the virological tests revealed a genotype D hepatitis B envelope antigen (HBeAg) positive CHB with a high viremia (HBV-DNA), mild elevation of ALT (50 IU/ml) and an unexpected low titer of HBsAb (26 mIU/ml, with a protective value above 100 mIU/ml). Anti-hepatitis D and C virus antibodies were negative.

A serological testing performed three years before was diagnostic for HBeAg negative CHB with moderate elevation of ALT (520 IU/ml), medium-low level of HBV-DNA, and absence of HBsAb, suggesting a subsequent seroreversion from HBeAg-negativity/anti-HBe positivity to HBeAg positivity. Till that time, the patient had neither received antiviral drugs nor indication for repeating virological or liver tests.

A liver biopsy was performed showing moderate necroinflammatory activity and bridging fibrosis (Stage 4/6 according to *Ishak's Score*) [7]. Because the fibrotic evolution, in spite of the genotype D of HBV and the immune-tolerance phase of CHB, antiviral treatment with recombinant IFN-alpha-2b (IntronA®) was began at the dose of 10 MU three times a week. This option was also supported by clinical guidelines at that time, with an expected anti-HBe seroconversion rate of about 37% [8]. During treatment, the patient was monitored by three-monthly blood tests and clinical visits (in Figure 1 six-monthly blood tests are reported).

After six months of interferon treatment, neither a biochemical response nor a virological response were observed, while HBsAb titer raised to 235 mIU/ml (Figure 1). Since consistent data about adefovir dipivoxil (ADV) were already available, the patient was started with ADV (10 mg/day) [9]. Although, after six months, a partial virological response was observed (decrease in HBV DNA of more than 1 log₁₀/ml), after twelve months, a virological breakthrough was detected (increase in HBV DNA level of more than 1 log₁₀/ml). We were at the beginning of 2007 and the novel carbocyclic analogue of 2' deoxyguanosine Entecavir (ETV) had been just licensed, showing outstanding results [10]. Moreover, due to the persistence of HBsAb with high viremia, mutations in the HBsAg region were investigated. Conventional sequencing and UDPS of the polymerase (pol) region of HBV to identify the presence of mutations in the HBsAg ORF associated to mutations on the overlapping rt ORF, possibly present in minor components of viral quasispecies, were performed. The methods for UDPS and elaboration of sequence data have been previously described [6]. Some mutations in both rt and HBsAg ORFs were detected (with frequency varying from 10 to 96%), but none of these are known to be associated with resistance to current HBV-specific NAs or to be involved in immunological escape (Additional file 1: Table S1).

Based on these results, ETV (0.5 mg/day) was added to ADV and, three months later, HBV-DNA levels became undetectable. On July 2009, after twelve months of ADV and ETV combination therapy, due to an apparent new virological breakthrough (Figure 1), ADV was replaced with the novel nucleotide analogue tenofovir (TDF). After three months of TDF (245 mg/day) plus ETV (0.5 mg/day) combination, virological suppression was achieved. Surprisingly, on January 2011, i.e. six months later, anti-HBe seroconversion and HBsAg loss were observed, while the titer of HBsAb persisted. Anti-HBe seroconversion and HBsAg loss were confirmed after six months, when antiviral agents were discontinued. Unfortunately, the patient died on September 2011 because of a ruptured thoracic aortic aneurysm, so further follow-up visits are not available to verify the sustained off-treatment virological response.



This is, to our knowledge, the first case described of a CHB with HBsAg/HBsAb positivity, wild type for mutations clinically relevant in HBsAg and rt ORFs, successfully treated with a combination of NAs.

Consent

Written informed consent was obtained from the patient for publication of this Case report and any accompanying images. A copy of the written consent is available for review by the Editor of this journal.

Additional file

Additional file 1: Mutation frequency in rt and HBsAg ORFs by UDPS.

Abbreviations

HBsAg: Hepatitis B surface Antigen; HBsAb: Anti HBsAg antibodies; CHB: Chronic hepatitis B; HBV: Hepatitis B Virus; ORFs: Overlapping open reading frames; Rt: Reverse transcriptase; NAs: Nucleot(s)ide analogues; HBeAg: Hepatitis B envelope Antigen; UDPS: Ultra-deep pyro-sequencing; NGS: Next generation sequencing; IFN: Interferon; ADV: Adefovir dipivoxil; ETV: Entecavir; Pol: Polymerase; TDF: Tenofovir; LLQ: Lower limit of quantification.

Competing interests

The authors declare that they have no competing interest.

Authors' contributions

GG conceived the clinical case and wrote the manuscript; AD helped to draft the manuscript and designed Figure 1 and Additional file 1: Table S1; UV helped to draft the manuscript and revised the written English; PG, DV, MCS, CD and AP helped to draft the manuscript and revised it critically for important intellectual content; DV and MCS performed the ultra-deep pyro-sequencing of the polymerase region of HBV; all authors read and approved the final manuscript.

Acknowledgements

Capobianchi Maria Rosaria, MD, PhD; Elisabetta Riva MD, PhD; Carlo Federico Perno, MD, PhD; Valentina Svicher MD.
Funding: Italian Ministry of Health (Ricerca Corrente).

Author details

¹Clinical Medicine and Hepatology Unit, Campus Bio Medico University of Rome, 200, Alvaro Del Portillo Street, Rome 00128, Italy. ²National Institute for Infectious Diseases "Lazzaro Spallanzani", 292 Portuense Street, Rome 00149, Italy.

Received: 29 January 2014 Accepted: 7 May 2014

Published: 17 May 2014

References

- Pungpapong S, Kim WR, Poterucha JJ: **Natural history of hepatitis B virus infection: an update for clinicians.** *Mayo Clin Proc Mayo Clin* 2007, **82**:967–975.
- Tabor E, Gerety RJ, Smallwood LA, Barker LF: **Coincident hepatitis B surface antigen and antibodies of different subtypes in human serum.** *J Immunol Baltim Md 1950 1977*, **118**:369–370.
- Lada O, Benhamou Y, Poynard T, Thibault V: **Coexistence of hepatitis B surface antigen (HBs Ag) and anti-HBs antibodies in chronic hepatitis B virus carriers: influence of "a" determinant variants.** *J Virol* 2006, **80**:2968–2975.
- Torresi J: **The virological and clinical significance of mutations in the overlapping envelope and polymerase genes of hepatitis B virus.** *J Clin Virol Off Publ Pan Am Soc Clin Virol* 2002, **25**:97–106.
- Hussain M, Lok AS: **Mutations in the hepatitis B virus polymerase gene associated with antiviral treatment for hepatitis B.** *J Viral Hepat* 1999, **6**:183–194.
- Solmone M, Vincenti D, Prosperi MCF, Bruselles A, Ippolito G, Capobianchi MR: **Use of massively parallel ultradeep pyrosequencing to characterize the genetic diversity of hepatitis B virus in drug-resistant and drug-naïve patients and to detect minor variants in reverse transcriptase and hepatitis B S antigen.** *J Virol* 2009, **83**:1718–1726.
- Ishak K, Baptista A, Bianchi L, Callea F, De Groote J, Gudat F, Denk H, Desmet V, Korb G, MacSween RN: **Histological grading and staging of chronic hepatitis.** *J Hepatol* 1995, **22**:696–699.

- Wong DK, Cheung AM, O'Rourke K, Naylor CD, Detsky AS, Heathcote J: **Effect of alpha-interferon treatment in patients with hepatitis B e antigen-positive chronic hepatitis B. A meta-analysis** *Ann Intern Med* 1993, **119**:312–323.
- Marcellin P, Chang T-T, Lim SG, Tong MJ, Sievert W, Shiffman ML, Jeffers L, Goodman Z, Wulfsohn MS, Xiong S, Fry J, Brosgart CL: **Adefovir Dipivoxil 437 Study Group: Adefovir dipivoxil for the treatment of hepatitis B e antigen-positive chronic hepatitis B.** *N Engl J Med* 2003, **348**:808–816.
- Dimou E, Papadimitropoulos V, Hadziyannis SJ: **The role of entecavir in the treatment of chronic hepatitis B.** *Ther Clin Risk Manag* 2007, **3**:1077–1086.
- Zhang J-M, Xu Y, Wang X-Y, Yin Y-K, Wu X-H, Weng X-H, Lu M: **Coexistence of hepatitis B surface antigen (HBsAg) and heterologous subtype-specific antibodies to HBsAg among patients with chronic hepatitis B virus infection.** *Clin Infect Dis Off Publ Infect Dis Soc Am* 2007, **44**:1161–1169.
- Colson P, Borentain P, Motte A, Henry M, Moal V, Botta-Fridlund D, Tamalet C, G  rolami R: **Clinical and virological significance of the co-existence of HBsAg and anti-HBs antibodies in hepatitis B chronic carriers.** *Virology* 2007, **367**:30–40.
- Hsu HY, Chang MH, Liaw SH, Ni YH, Chen HL: **Changes of hepatitis B surface antigen variants in carrier children before and after universal vaccination in Taiwan.** *Hepatol Baltim Md* 1999, **30**:1312–1317.
- Frider B, Alessio A, Pozzati M, Cuestas ML, Mathet VL, Oubi  a JR: **Successful interferon treatment in a patient chronically infected with hepatitis B virus carrying unusual S- (and P-) mutants in the presence of anti-HBs antibodies.** *Liver Int Off J Int Assoc Study Liver* 2007, **27**:727–730.
- Lok AS, Trinh H, Carosi G, Akarca US, Gadano A, Habersetzer F, Sievert W, Wong D, Lovegren M, Cohen D, Llamoso C: **Efficacy of entecavir with or without tenofovir disoproxil fumarate for nucleos(t)ide-na  ve patients with chronic hepatitis B.** *Gastroenterology* 2012, **143**:619–628.e1.
- Brunetto MR, Giarin MM, Oliveri F, Chiaberge E, Baldi M, Alfarano A, Serra A, Saracco G, Verme G, Will H: **Wild-type and e antigen-minus hepatitis B viruses and course of chronic hepatitis.** *Proc Natl Acad Sci U S A* 1991, **88**:4186–4190.

doi:10.1186/1471-230X-14-94

Cite this article as: Galati et al.: Coexistence of HBsAg and HBsAb in a difficult-to-treat chronic hepatitis B: loss of HBsAg with entecavir plus tenofovir combination. *BMC Gastroenterology* 2014 **14**:94.

Submit your next manuscript to BioMed Central and take full advantage of:

- **Convenient online submission**
- **Thorough peer review**
- **No space constraints or color figure charges**
- **Immediate publication on acceptance**
- **Inclusion in PubMed, CAS, Scopus and Google Scholar**
- **Research which is freely available for redistribution**

Submit your manuscript at
www.biomedcentral.com/submit

